



REC'D 13 JAN 2004

WIPO

PCT

Kongeriget Danmark

Patent application No.: PA 2002 01896

Date of filing: 11 December 2002

Applicant: Ferrosan A/S
(Name and address) Sydmarken 5
DK-2860 Søborg
Denmark

Title: Gelatin materials as swabs

IPC: -

This is to certify that the attached documents are exact copies of the above mentioned patent application as originally filed.

PRIORITY DOCUMENT
SUBMITTED OR TRANSMITTED IN
COMPLIANCE WITH
RULE 17.1(a) OR (b)



Patent- og Varemærkestyrelsen
Økonomi- og Erhvervsministeriet

29 December 2003

Pia Høybye-Olsen
Pia Høybye-Olsen



PATENT- OG VAREMÆRKESTYRELSEN

1

Patent- og
Varemærkestyrelsen

11 DEC. 2002

Modtaget

GELATIN MATERIALS AS SWABS

Field of the Invention

- 5 A gelatin-based sponge or powder are used in the collection of targets such as microbiological and mammalian cells, detergents, nucleotides and nucleic acids, from an array of collection media, such as solid work surfaces and mammalian tissues.

Background of the Invention

10

Gelatin-based sponges have been used as haemostatic agents in surgical procedures.

Dean Jr. (US 4,997,753; US, 4,863,856 and US 4,861,714) discloses the use of weighted collagen microsponges for immobilising bioactive materials.

15

EP 0 702 081 discloses a matrix for tissue culturing comprising two kinds of sponges.

US 5,462, 860 discloses a culture medium for rapid growth and detection of microbes.

20

Conventional methods of sampling an area for a target area comprise the use of cotton swabs. Sample recovery from the surface is limited due to the low recovery from the surface area to the swab and then low transfer from the swab to a culturing medium. The level of recovery of micro-organisms from surfaces is critical when complying with USP/NF guidelines and EU-GMP guidelines. The present invention addresses this problem and provides for a dramatically

25

improved device for sampling an area.

Summary of the Invention

- 30 A gelatin-based sponge has been found to be useful in the collection of a variety of materials, such as microbes and mammalian cells, but also mammalian tissue and various molecules, including nucleotides, from an array of collection media. The sponge can be used for instance for sampling and culturing purposes. The present inventors have surprisingly found that the sponge has a dramatically higher recovery level of the sampled material than conventional methods. Moreover, the target material bound to the sponge can be transferred from the
- 35 sponge to another medium by an array of methods.

A first object of the invention relates to a device comprising

i) a gelatin-based sponge; and

ii) a support

40

wherein said sponge comes into contact with or is attached to said support.

The device or the gelatin-based sponge may be used for an array of applications related to the high recovery of targets from the sponge. Accordingly, a further object of the invention relates to the use of a gelatin-based sponge for collection of a target from a collection medium comprising making contact between the gelatin-based sponge and the medium. Furthermore, 5 an object of the invention relates to a method of lowering the amount of a target in a sample area comprising making contact between a gelatin-based sponge and at least a portion of said sample area, such that the target adheres to the sponge. An important and further utilisation of the surprising properties of the sponge relate to a method of qualitatively or quantitatively sampling an area for content of a target comprising the use of a gelatin-based sponge and 10 steps of i) wet sampling using the gelatin-based sponge; and/or ii) dry-sampling using the gelatin-based sponge. A similarly important aspect of the invention relates to a method for culturing micro-organisms or mammalian cells comprising adhering the cells to a gelatin-based sponge and culturing the cells in a growth medium.

15 Description of the Invention

In the present context, a target refers to any species which binds to the gelatin-based sponge of the invention. A collection medium refers to any medium from which said targets may be collected. The term "transfer medium" is intended to mean a medium to which the collected 20 target is transferred.

In the present context, the term "recovery" is intended to mean the overall recovery yield of a target from a collection medium to transfer medium. Thus, recovery comprises i) the collection yield of a target to the collagen-based sponge of the present invention, as well as ii) the 25 transfer yield from the collagen-based sponge to a transfer medium.

A first transfer medium is, in the present context, considered to be a medium used in the collection of a target from the sponge. Suitable examples of a first transfer medium include an enzymatic solution, or a suitable washing agent to mechanically or chemically remove the target 30 from the sponge into the medium, such as a liquid medium. A second transfer medium is considered to be represented by a second medium into which the first medium or the sample from the first medium, which includes the collected target, is transferred.

In the present context, a micro-organism is considered to be any organism selected from the 35 group consisting of bacteria, bacterial spores, viruses, archaea, yeast and fungi.

A dispersion agent is, in the present context, considered to be any liquid agent into which targets may be dispersed following collection.

40 A neutral diluent is considered to be a liquid which is neutral in the context of the assay process, i.e. it does not interfere with the diagnostic assay being performed. In this context, therefore, a neutral diluent is not necessarily, but can be, a diluent with neutral pH. Water, aqueous buffer solutions are suitable examples of neutral diluents.

Ringer's solution refers to an aqueous solution comprising distilled water and sodium chloride, potassium chloride, and calcium chloride at roughly the same concentrations as their occurrence in body fluids.

- 5 A "semi-solid surface" is a surface which is not strictly solid in nature, such as mammalian tissue or any other natural and/or synthetic tissue. Suitable examples of semi-solid surfaces are mammalian skin or other surfaces covered by connective tissue, such as surfaces of mammalian organs.
- 10 In the present context, a "detergent" can be any natural or synthetic, organic or inorganic, compound or a mixture of compound used for cleaning purposes, such as for the removal of impurities or contaminants from a surface.

15 The term "handle" in the present context is to be considered to relate to any device that can be used for gripping, and should not be construed to be limited to devices specially designed to act as a handle. By way of an example, a stick attached to the gelatin-based sponge of the invention can thus be used to handle the sponge, and is therefore considered to represent a handle. Moreover, a tweezer or tong, which only temporarily connect to the sponge, is to be considered a handle.

20 A swab is in the present context considered to be any material used for applying or removing material from an area or a surface. Swabbing refers to the method of applying or removing material from an area or surface using said swabs.

- 25 The present invention relates to gelatin-based sponges and their properties with respect to binding of targets, such as micro-organisms and mammalian cells, as well as molecular species such as nucleotides, which are suitable for binding to the gelatin-based sponges. The binding properties of the aforementioned targets are useful for the collection of said targets from various media. Once the target is collected, it may be optionally be released from the sponge by
- 30 mechanical, enzymatic, or chemical processes.

Collection media

- In one embodiment, the medium is a solid surface from which targets may be collected. In a
- 35 second embodiment, the medium is a liquid, from which targets may also be collected. In such an embodiment, the high water absorption capacity of the gelatin-based sponge is a useful characteristic, as it allows collection of large volumes of water. The liquid medium may be located on a surface, for example in cavities on the surface. The sponge therefore can be adapted to be useful for the collection of targets from a wide variety of sources, such as
- 40 manufacturing devices in food manufacturing, processing plants for meat and/or fish products, medical devices, as well as the management and cleaning of wounds, such as surgical wounds.

It is furthermore envisaged that the gelatin-based sponge may be adapted for use in collecting targets from other types of samples of liquid and/or gaseous nature.

- 45 In principle, collection may be from a solid surface irrespective of the material from which the surface is comprised, such as natural or synthetic surfaces, of organic or inorganic material.

Furthermore, targets may be collected from semi-solid surfaces, such as mammalian surfaces and mammalian tissue, including mammalian skin and the surface of mammalian organs.

The micro-organism may be selected from the group comprising bacteria, archaea, bacterial spores, yeast, fungi and viruses.

The mammalian cell may be selected from the group consisting of cells from blood plasma, leukocytes, erythrocytes, thrombocytes, but also other mammalian cells such as skin cells or any other type of mammalian cells that may be useful to collect for various diagnostic and/or cleaning purposes

Properties of the gelatin-based sponge

The gelatin-based sponge is characterised by its physical properties, which in particular may be described by the gelatin composition, pore size, reconfirmation rate, water absorption and digestibility of the sponge.

Without being bound to a particular theory, the pore size of the sponge is considered to influence, at least in part, the ability of the sponge to collect and transfer the targets of the invention. Moreover, the pore size influences the density of the sponge, and also has an effect on its physical characteristics, such as reconfirmation rates and water absorption.

As stated, the sponge is gelatin-based. The gelatin-based sponge material is preferentially comprised of at least 50% gelatin, such as at least 60%, such as at least 70%, typically at least 75% gelatin, preferably at least 80%, more preferably at least 85% gelatin, such as at least 90% gelatin, suitably at least 95% gelatin, most preferably selected from at least 96%, 97%, 98%, 99% gelatin, based on the dried weight of the sponge.

In one embodiment of the invention, the gelatin-based sponge is based on gelatin of porcine origin. It is envisaged that the invention may be adapted to include gelatin with other origins, such as gelatine of bovine, or any other mammal, including marine mammal, fish or crayfish or vegetable origin, and including gelatin of any other origin, such as of gelatin of organic origin, or synthetic or semi-synthetic origin.

Reconfirmation rates represent a measure of the elasticity of the gelatin-based sponge. In one embodiment of the invention, the gelatin-based sponge has a reconfirmation rate of no more than 10 seconds, and typically no more than 5 seconds. However, it is envisaged that sponges based on gelatin from various sources will have a wider range of reconfirmation. The reconfirmation rate is typically determined by a method based on the rate at which the sponge regains its original size and shape, as described in Example 1.

The gelatin-based sponge of the invention typically has a water absorption capacity which is in the range of at least 30g/g and more typically at least 40g/g, as determined in Example 3. It is however envisaged that the water absorption capacity of sponges based on gelatin from various sources may be in a wider range of at least 5g/g, such as at least 10g/g or at least 20g/g. The determination of water absorption is typically performed according to USP standards.

The inventors have found that the gelatin-based sponge binds micro-organisms tightly but reversibly. Thus, the sponge may be used for sampling for the presence of micro-organisms from a sample, such as from a surface. In such an embodiment, the micro-organism can be transferred from the sponge by the methods of the invention. The thus collected micro-organisms may optionally be furthermore cultivated, which may find uses for specific purposes such as detailed characterization of said micro-organism. Alternatively, the sponge may be used for quantitative removal of micro-organisms from a sample, such as a surface. In such an embodiment, anti-microbial or disinfecting agents may optionally be suitably incorporated into the gelatin-based sponge.

It is furthermore envisaged that the gelatin-based sponge may be adapted for use in collecting other types of cells in addition to micro-organisms, such as mammalian cells. Such an embodiment may be realized by, e.g. collection from mammalian surfaces, such as from mammalian skin or the surface of any mammalian organ. Furthermore, the gelatin-based sponge may be adapted for internal use, such as during surgical operations on a mammal, and may in such embodiments be used to collect targets from wounds or from internal organs of a mammal, as well as surgical equipment and specialized furniture, walls or floors in a health clinic or a hospital, such as in surgical rooms, or in any other facility used to conduct or perform surgical procedures.

The inventors have furthermore found that the gelatin-based sponge binds certain molecules, such as purine- or pyrimidine-based nucleotides or nucleic acids, preferentially ATP, in a reversible fashion. This finding may find a useful application in that bacterial numbers have been estimated in foods by measuring the amount of bacterial adenosine triphosphate (ATP). It is envisaged that the gelatin-based sponge may more generally be adapted to collect a variety of molecular species, and thus the gelatin-based sponge may find general use for collection of certain molecules.

In one embodiment, such molecules are purine- or pyrimidine-based nucleotides, such as ATP. In another embodiment, such molecules are nucleic acids. In yet another embodiment, the molecules are detergents, which may conveniently be collected from a sample, such as a surface. In this context, detergents can be any natural or synthetic, organic or inorganic, compound or a mixture of compound used for cleaning purposes, such as for the removal of impurities or contaminants from a surface.

Furthermore, the gelatin-based sponge may be adapted to collect a mixture of micro-organisms, mammalian cells and/or molecules simultaneously from a sample. In certain embodiments, it may be useful to configure the gelatin-based sponge such that one particular type of target is collected.

The desirable property of the gelatine-based sponge in that it is capable of adhering to a target such as micro-organisms or mammalian cells as well as molecular species, especially organic molecules such as nucleotides, make a variety of useful applications possible.

In a suitable embodiment of the invention, the gelatin-based sponge is adapted so that it comes into contact with or is attached to a support. The purpose of a support can be that of providing a way of handling the gelatin-based sponge without touching the sponge material itself, and thus avoiding contamination. This may be especially useful for embodiments in which the
5 sponge is ideally pre-sterilized, i.e. in embodiments for the collection or analysis of micro-organisms or mammalian cells. A support may also facilitate the use of the gelatin-based sponge, and allow convenient collection of targets from various samples. A support can furthermore be invaluable for the collection of targets from samples that may be difficult to reach or for other applications in which the sponge is difficult to manipulate without a support.

10

The support may be made of any suitable material for the particular use such as wood, natural or synthetic polymeric material, including plastics and rubber materials, or any other organic or inorganic material suitable for the particular embodiment.

- 15 The support may be of a wide variety, such as conveniently in the form of a handle. The handle may be short, such as in the range of about 1 cm to about 30 cm, such as about 3 cm to about 20 cm, preferably about 5 cm to about 15 cm. For certain applications, a handle may suitably be considerably larger, such as in the range of about 30 cm to several meters in length. The handle may be of any shape convenient for the particular embodiment, but is typically
20 elongated, optionally bent, with the gelatin-based sponge attached at one end of the handle, while the other end of the handle is used for gripping and otherwise applying the gelatin-based sponge

- In one suitable embodiment, in which the gelatin-based sponge is attached to a support in the
25 form of a handle, the gelatin-based sponge has an oval or otherwise elongated shaped sponge positioned at the end of a stick, which serves the function of a handle. In another embodiment, the sponge is attached to a solid support, such as a circular or rectangular support, which in turn is positioned on the end of a handle or stick. It should however be appreciated that the sponge may be adapted to be attached to a support in the form of a stick or handle in multiple
30 different embodiments, suitably adapted for any given use of the gelatin-based sponge.

- In another embodiment, the support is optionally in the form of a coating. The coating may be comprised of any suitable material, e.g. polymeric material or plastic material, or any other material suitably used to provide a coating. In a suitable embodiment, the coating is applied to
35 one side of a gelatin-based sponge of the invention which has been adapted to be in a flat shape.

- In yet another embodiment, the support is a solid material which has a suitably adapted shape, such as the shape of a disc, cube, sphere or a block. In such embodiments, the gelatin-based
40 sponge is preferably attached to one side of the support, but may be attached to several or all surfaces of the support in preferred embodiments.

- An embodiment in which the sponge is of a cubical shape may be particularly useful is an
embodiment for collection of liquid samples or collection from surfaces comprising a liquid
45 coating. In such embodiments, the sponge may suitably be attached to a support, which may optionally be in the shape of a handle, or suitably attached to a stick or a handle.

It should be appreciated, that due to the nature of the gelatin-based sponge material, it may be adapted to a shape for any particular use. Thus, the dimensions and shape of the embodiments wherein the invention is realised by mounting the gelatin-based sponge onto a support, will be adjusted as ideally suited for their uses.

5

The sponge may be attached to the support by any conventional method known to those skilled in the art. The nature of such embodiments will depend on the particular shape of the embodiment and its intended use.

- 10 It is furthermore envisaged that the support may be in the form of a porous container, such as a crucible or otherwise suitable material shaped such that it encloses the gelatin-based sponge, while allowing liquid and targets to pass through the enclosing material. In such embodiments, the encased gelatin-based sponge can be used to collect samples from a liquid. Such collection may suitably be performed by immersing the encased sponge into the liquid medium, thus
15 allowing targets in the medium to come into contact with and bind to the gelatin-based sponge.

In embodiments of the gelatin-based sponge in which it is adapted to collect gaseous targets, it is preferably adapted such that the surface area and/or pore size of the sponge is maximised. Such embodiments may be realised in accordance with any of the embodiments disclosed.

- 20 above, since the nature of binding of gaseous targets follows the same principles as that of targets in a liquid medium. Gaseous targets may be any molecular species, which is gaseous at the temperature applied during collection, or is a liquid or a solid compound which has a vapour pressure high enough so that the target may be collected. A gaseous target is in this context considered to include targets which are solid in nature, including micro-organisms and
25 mammalian cells, but either are trapped in liquid droplets or microdroplets or form particles that may be carried by gases, such as ambient atmosphere.

- In yet another embodiment of the invention, the gelatin-based sponge is in the form of a powder. The gelatin material of the sponge may form a powder by using conventional
30 techniques known to those skilled in the art. The particle size of such powder can be adapted to the specific use, and is conveniently in the range of about 1 μm to about 1 mm, such as about 5 μm to about 0.5 mm, such as about 10 μm to about 0.1 mm. Such powder may be adapted for use according to the methods of the invention by using the powder directly. Alternatively, the powder is incorporated into a suitable binding material to which it will adhere, so that it may be
35 used for collection of targets. Suitable binding materials include commercially available materials and tools known to those skilled in the art. One example is swabs, such as cotton swabs, but other convenient materials may also be used, optionally positioned on supports, such as supports in the form of a stick.

- 40 The gelatin powder may also be enclosed by a crucible or other suitable material shaped such that it encloses the powdered gelatin-based sponge, while allowing liquid and targets to pass through the enclosing material. Such embodiments may in particular be useful for collection from liquid media. In yet another embodiment, the enclosing material is permeable to gaseous targets, and the collection of targets is realized by placing the encased powdered gelatin
45 material in an environment containing said gaseous targets.

Alternatively, the powdered gelatin-based sponge may be used without a support. In such embodiments, the powder may be applied directly to a sample for collection of targets, where the sample is suitably a liquid. The gelatin powder may in such embodiments be recovered by filtration, centrifugation or by other means known in the art.

5

Methods used for collecting targets from a surface using the gelatin-based sponge of the present invention include techniques such as swab techniques and count-tact techniques. Swab techniques involve in principle a mechanical swiping or swabbing of a target, such as a surface, and are well known to those skilled in the art. Count-tact techniques involve the use of a
10 specially designed instrument, as disclosed in Example 6.

In one embodiment of the invention, the gelatin-based sponge, optionally attached to a support, is used to collect targets from a sample. The target may be selected from the group comprising a micro-organism, mammalian cell, but may also be mammalian tissue, or
15 alternatively a molecular species, such as a nucleic acid or a purine- or pyrimidine-based nucleotide, preferentially ATP. The micro-organism may be selected from the group consisting of bacteria, archaea, bacterial spores, yeast, fungus or a virus. The mammalian cell can be of any mammalian cell type, but may in particular be selected from the group consisting of cells from blood plasma, leukocytes, erythrocytes, thrombocytes, epithelial cells, skin cells or any
20 other mammalian cell type that may be useful to collect for various diagnostic and/or cleaning purposes.

In a preferred embodiment, targets are collected from a surface using a combination of wet-sampling and dry-sampling. In this context, wet-sampling is considered to comprise the use of
25 a gelatin-based sponge of the invention, which optionally has been attached to a support and has been pre-wetted with a suitable neutral diluent or a dispersing agent. Such a diluent or dispersion agent serves the purpose of facilitating the recovery of the targets from the surface, while not interfering with the assay. The diluent and dispersion agents may be any suitable aqueous solution, optionally including salts or other agents not toxic or otherwise chemically or
30 biologically harmful for the targets to be collected, such as saline, saline peptone, buffered saline peptone, Ringer solution and an organic or inorganic buffer, optionally containing inorganic salts. The diluent or dispersion agent may also optionally contain growth media suitable for the micro-organism or mammalian cell type being collected, for assays directed towards such targets. The sponge may also optionally be presterilized. Collection of targets is
35 realized by swiping the surface with at least one such pre-wetted sponge, followed by swiping of said surface with at least one dry gelatin-based sponge. The purpose of the dry swiping is to recover as much as possible of remaining liquid and target from the surface.

Collection of targets may, in other embodiments, be realised by wet-sampling or dry-sampling
40 alone. The choice of method to use will vary depending on the sample type to be assayed and the target type to be collected.

Targets, i.e. micro-organisms, mammalian cells and/or molecules collected in the gelatin-based sponge may optionally be transferred from the sponge. Transfer of such collected targets
45 comprises removing or unbinding such targets from the sponge into a suitable medium. In a preferred embodiment, this is accomplished by placing the swabs in a medium comprising a solution capable of digesting the gelatin-based sponge. Digestion of the sponge may be

realised by chemical and/or enzymatic methods, preferably using enzymes such as proteases, more preferably using proteases such as alcalase or pepsin. Digestion by chemical means may comprise using mineral or carboxylic acids, or bases, in appropriate concentration not to denature the target. In one embodiment, digestion comprises using a mixture of at least one
5 enzyme, and may optionally include a mineral acid or a base, and optionally inorganic salts as well as organic or inorganic buffering agents. The temperature suitable for recovery of targets from the sponge will be highly dependent on the method used. In embodiments wherein transfer is realised using enzymes, the experimental temperature will be adjusted so as to maximise the efficiency of digestion for the particular enzyme in the context of the composition
10 of the digestion medium employed.

In general, transfer of targets from the gelatin-based sponge of the invention may be realised by any technique known in the art which releases said target from the sponge. Thus, this may occur by changes in conditions such as pH or temperature, or by adding salts, chaotropic
15 agents or organic solvents. Transfer from the sponge may also optionally include mechanical action, such as that generated by rubbing or shaking the sponge, or by other mechanical means facilitating the unbinding of targets from the sponge. Transfer from the sponge may furthermore be realized by washing of a target from the gelatin-based sponge.

20 It is envisaged that preferred embodiments of the invention will include digestion methods for releasing micro-organisms and/or mammalian cells from the sponge, since micro-organisms and mammalian cells are in general sensitive to changes in conditions. However, there may be exceptions to this, and in particular it is envisaged that for certain applications it will be useful to use changes in experimental conditions for recovery of certain microorganisms or
25 mammalian cells, in particular extremophiles, which are organisms adapted to withstand harsh conditions of pH, salt, temperature and/or organic solvents. Recovery of bound molecules to the sponge may be accomplished by any of the aforementioned techniques, depending on any given embodiment of the invention and the type of molecule bound.

30 Micro-organisms or mammalian cells recovered from the gelatin-based sponge may optionally be further isolated using membrane filtration, wherein the membrane filter has properties such that it allows solvent and small molecules to pass through the filter, while whole cells and micro-organisms do not. In a typical embodiment, such a filter has a pore size of less than 1 μm , such as less than 0.8 μm , such as less than 0.6 μm , more typically less than 0.45 μm , such
35 as less than 0.2 μm .

In another embodiment of the invention, the gelatin-based sponge is used to disinfect a sample, such as a surface. In such an embodiment, the combined action of the sponge, which acts to remove targets, in this case micro-organisms and/or mammalian cells, from a sample, and
40 optionally an anti-microbial or a disinfecting agent, facilitates the effective removal of said micro-organisms or mammalian cells from the sample, thus rendering it sterile. The sponge is in such embodiments preferably sterilised by methods known in the art, such as by heat and/or radiation, and optionally pre-treated with an anti-microbial or disinfecting agent. Such an agent is preferably a liquid, such as an alcohol, an aqueous solution comprising an alcohol or other
45 liquid agent which kills micro-organisms or mammalian cells, but may be any compound which facilitates the sterilisation procedure. Embodiments for sterilisation of samples may be realised by packaging individual gelatin-based sponges, optionally attached to a support and optionally

pretreated with a sterilising agent, individually into sealed packages, which are ideally intended for single use.

5 In another embodiment of the invention, viable micro-organisms or mammalian cells which have been trapped on the gelatin-based sponge can be cultured. Culturing in general requires contacting a micro-organism or mammalian cell and a suitable growth medium. The growth medium may in the form of a liquid; alternatively, it is in the form of solid agar. The growth medium is comprised of components well known to those skilled in the art. Realisation of said culturing can be accomplished by conventional techniques, including:

10

1. Contacting the gelatin based sponge, optionally attached to a support, and a liquid or a solid growth medium. In one such embodiment, a suitably shaped gelatin-based sponge positioned at the end of a stick is used to collect targets from a surface, and subsequently allowed to come into contact with a growth medium, such that micro-organisms or mammalian cells collected by
15 the sponge are transferred to the growth medium.

2. Injecting a liquid growth medium, optionally agar-containing, into the gelatin sponge, thus providing conditions for *in situ* growth of the bound micro-organisms or mammalian cells in the gelatin-based sponge.

20

3. Transferring the gelatin-based sponge to a container with liquid growth medium, thus allowing for culturing of the entire population of bound micro-organisms or mammalian cells in said growth medium.

25 It should be appreciated that the culturing of collected micro-organisms or mammalian cells may be preceded by a step in which bound cells have been unbound or by other means released into a medium. In such embodiments, said medium is considered to be a first transfer medium. The medium can be any liquid suitable for the application, such as a neutral diluent, a dispersion agent, or a growth medium. Unbinding of bound micro-organisms or mammalian
30 cells may be realized by any of the methods described herein, including enzymatic and/or chemical digestion of the gelatin-based sponge, and may include mechanical transfer into said first transfer medium. The thus unbound micro-organisms or mammalian cells are subsequently transferred to a second transfer medium, which is ideally comprised of a liquid or solid growth medium. The transfer to a second transfer medium may be partial, i.e. a sample
35 from said first growth medium is transferred to said second transfer medium. Alternatively, the transfer is complete, i.e. the entire volume of the first transfer medium is transferred into the second transfer medium.

40 Culturing of micro-organisms or mammalian cells collected by the gelatin-based sponge of the invention may in particular be useful for further characterisation or production of the collected micro-organisms or mammalian cells. Preferred embodiments of the gelatin-based sponge may, for example, be used for qualitative determination of the microbiological and/or mammalian cell composition of a target population collected from a sample. In such
: embodiments, the sample may, for example, be a surface in a food production line, such as
: 45 meat or fish processing lines, or from other surfaces such as floors, walls, or equipment used in such processing lines. Alternatively, the sample may be a surface from equipment, specialized furniture or walls or floors from a health clinic or hospitals, such as surgical rooms. The sample

may also be collected from an open wound, from the surface of an internal organ or it may be comprised of any mammalian tissue. In principle, however, the gelatin-based sponge may be adapted for collecting and culturing targets from any sample, from which it is useful to determine the microbiological and/or mammalian cell content.

5

Examples

The following methods and examples illustrate how the gelatin-based sponge of the present invention may for specific uses be adapted for the collection and recovery of bacterial spores from a stainless steel surface. Recovery yields, i.e. the overall yields for the transfer from the surface to the sponge and the subsequent transfer from the sponge to a medium, are very high which is an illustration of the usefulness of the gelatin-based sponge for the collection and transfer of targets from a sample, such as a stainless steel surface.

15

These methods and examples should however only be understood as examples of useful embodiments of the present invention, and in no way limiting for its adaptation for other use.

Example 1. Determination of reconfirmation time of gelatin-based sponges

20

The purpose of this method is to determine the reconfirmation rate of a gelatin-based sponge. The method comprises soaking the sponge, and subsequently squeezing it. The appearance of the native shape of the sponge is monitored as a function of time, and the time that lapses until the sponge has reached its native shape is termed the reconfirmation time.

25

The method comprises the following steps:

1. Cut a suitable piece of absorbable gelatin-based sponge, approximately 1x1 cm, and thoroughly soak it in water at room temperature.

30

2. Remove the sample from the water, and squeeze it until it is flat and no more air bubbles or drops of water can be pressed out.

3. Place the sample in a beaker filled with water at room temperature and measure the time (in seconds) until the sample has gained its former size and shape.

35

4. Repeat the test twice and report the result as the average of three determinations.

Example 2. Determination of digestability of gelatin-based sponges using pepsin

40

Purpose: To determine the digestion time of a gelatin-based sponge by enzymatic means using pepsin.

Reagents used in the method:

45

Milli-Q-water

Pepsin(1:3000)

Hydrochloric acid dilute, Ph. Eur.

12

Pepsin solution 1 %:

Accurately weigh 20.0 g of pepsin and dissolve in 100 ml hydrochloric acid dilute, using a 2000 ml volumetric flask. Add water to volume and mix.

To be prepared daily

5

Apparatus:

Metallic filament basket, 6 cm Ø, 1 mm openings.

Thermostated water bath

10

Method:

1. Transfer 100 ml of the pepsin solution (1%) to a 250 ml beaker. Add a magnet and place the beaker in a water bath, previously warmed to 37 °C on a magnetic stirrer.

15

2. Cut off a piece of absorbable gelatin sponge weighing 50 ± 5 mg, and place it in a beaker of water. Knead gently between the fingers until the gelatine sponge is thoroughly wet, and until all the air has been removed, taking care not to break the tissue.

20

3. Lift from water and squeeze gently to remove any excess water.

4. Place the wetted sample in a metallic filament basket in the beaker, and start the timer.

5. Watch until the piece of absorbable gelatin sponge is entirely dissolved, stop the timer, and note the consumption of time.

25

6. Repeat the test twice - a total of three times and calculate average.

30 Alternative method for a gelatin powder

1. Transfer 100 ml of the pepsin solution (1%) to a 250 ml beaker. Add a magnet and place the beaker in a water bath, previously warmed to 37°C on a magnetic stirrer.

35

2. Prepare a sample weighing out approximately $50 \text{ mg} \pm 5 \text{ mg}$.

3. Place the sample directly in the beaker.

4. Watch until the absorbable gelatin powder is entirely dissolved, stop the timer, and note the consumption of time.

40

5. Repeat the test twice - a total of three times and calculate average of three determinations.

Example 3. Determination of water absorption of gelatin-based sponges

Purpose: To determine the amount of water that a gelatin-based sponge can absorb. The sponge is expected to absorb several times its own weight of water on a weight to weight basis.

- 5 Method: According to USP Method "Absorbable Gelatin Sponge: Water absorption". A total of 6 determinations on 6 different pieces of gelatin-based sponge are performed.

Example 4. Size measurements of a gelatin-based sponge

10

Purpose: The dimensions weight, height, length, width, centre hole and diameter of the gelatin sponge are measured on 6 samples; one series of measurements is performed for each of the 6 samples. The average of the 6 measurements is reported. The density is calculated.

- 15 Apparatus: A caliper, Mitutoyo 500-Series or similar.

A ruler specifically made for the determination of length and width of absorbable gelatin sponge film.

- 20 Balance, Mettler AK 160 or a balance with similar accuracy.

Method and calculations: The wanted measurements are made with a caliper or a ruler. The average of 6 measurements performed on 6 samples is reported (in mm).

- 25 The weight of the sponge is measured. The average of 6 measurements performed on 6 samples is reported (in 0.001 g).

The density of the absorbable gelatin sponge, anal is calculated in the following way:

30
$$\frac{4 * weight}{\pi * length * (D^2 - d^2)} = density, \quad mg / mm^3$$

where

D = diameter of the sponge.

- 35 d = diameter of the pore in the sponge.

Example 5. Sampling protocol from a stainless steel surface using wet sampling and dry sampling.

40

Purpose: To sample a stainless steel surface using a combination of wet sampling and dry sampling.

14

Materials:

A gelatin-based sponge.

Saltwater-peptone solution

Alcalase solution

5 A stainless steel surface with a 24 cm² contact area.

Stomacher bags

Method:

10 In this procedure, gloves should be used.

1. Swab the stainless steel sheets horizontally from left to right covering the 24 cm² contact area using a gelatin sponge swab moistened with saltwater-peptone solution.

15 2. Swap the stainless steel sheets vertically from top to bottom covering the 24 cm² contact area using a gelatin sponge swab moistened with saltwater-peptone solution.

3. Swab the stainless steel sheets horizontally from left to right covering the 24 cm² contact area using a dry gelatin sponge swab.

20

4. Swap the stainless steel sheets vertically from top to bottom covering the 24 cm² contact area using a dry gelatin sponge swab.

5. Place the swabs in stomacher bags.

25

6. Add the digesting solution to each stomacher bag.

7. Homogenize the stomacher bags using a stomacher.

30 8. Incubate the stomacher bags at 36°C until the swabs have been dissolved.

9. Extraction of the spores is carried out using membrane filtration.

35 **Example 6. Swabbing by Count-Tact applicator.**

Introduction. The Count-Tact applicator standardizes surface testing, by applying a uniform pressure of 500 ± 50 g for 10 ± 1 seconds (draft European Standard: CEN/TC 243).

40 **Description.** The applicator is composed of two plastic elements:

- the base, which holds the Count-Tact plate in position, consisting of a push-button device mounted on a calibrated spring;
- a unit clipped onto the base, containing the electronic timer, the audible beep mechanism and batteries.

45

15

The method involves the following steps:

1. Slide the Count-Tact plate into position (lid facing outwards) under the two transparent clips fixed on the bottom of the base.
2. Remove the lid from the Count-Tact plate.
- 5 3. Hold the applicator in position against the surface where the sample is to be taken, without moving it (check that the surface is not damp).
4. After 10 seconds of contact with the surface, the audible beep will sound. Remove the applicator from the surface. Place the lid back on the plate, and remove the plate from the applicator (do not forget to clean the surface where the sample was taken in order to remove
- 10 any possible traces of agar).
5. For incubation of the plates, refer to the Count-Tact technical sheet and the Count-Tact irradiated sheet.

15 Example 7. Validation Protocol from polished stainless steel.

Introduction. This protocol describes the validation of a method that is used for the microbiological sampling from polished stainless steel, which has been cleaned with Isopropanol 70%.

20

The validation protocol is carried out in order to define the accuracy and precision of the method, and the recovery yield of micro-organisms from the type of material used in this test, using the described method of sampling.

- 25 This current test does apply whenever microbiological sampling from polished stainless steel cleaned with isopropanol is carried out in accordance with the current described sampling procedures.

Validation parameters.

30 Selectivity

Bacillus spores are used as markers of microbiological activity on the stainless steel. Because the spores are incubated at 55°C, it is considered unlikely that any microbiological contamination will influence the results during the validation test. Thus, selectivity is not relevant for this test.

35

Precision (repeatability) and accuracy (recovery)

Assessed by replicate extraction and analysis of swab samples using the same analyst, same equipment and same reagents within the same day, using

2 concentration levels (5 and 25 spores)

40

1 surface material

2 wipings of each surface material (pooled to one sample for analysis)

Total of six replicates (n=6)

Therefore, a total of 12 samples will be analyzed.

45 Linearity

16

Established using a three-point calibration curve, covering ranges from 5 to 50 spores. Test is carried out in six replicates, using three concentration levels (5, 25 and 50 spores), and 2 wipings of each surface material, pooled for analysis.

5 Intermediate precision

Established by replicate analysis on different days, using same reagent batches, same equipment, different analysts. Demonstrated using 2 concentration levels, 5 and 25 spores, 1 surface material and two wipings (pooled for analysis), and six replicates by three analysts on three days.

10 Therefore, a total of 108 samples.

Equipment.

Incubators, autoclave, sterile Drigalski spates, sterile gloves, gelatin swabs, mixer, membrane filtration equipment, colony counter, sterile stomacher bags.

15

Materials.

Spore suspension (1×10^6 spores/0.1 mL ethanol) of *Bacillus stearothermophilus* spores in 40% ethanol.

20 Test samples of surface material (polished stainless steel).

Template for swabbing area ($6 \times 4 \text{ cm} = 24 \text{ cm}^2$)

Tryptic Soy Agar

Sterile Filter

Sterile 50 mL syringe

25 Disposable sterile pipette tips

Count-Tact applicator with contact plates containing TSA, Tween and Lecithin.

Chemicals

Saltwater-peptone solution

30 Ethanol 96%

Elga water

Alcalase solution

Preparation of Test Validation Solutions

35 Positive control

Class A: Spore suspensions containing 5 spores/313 μL is used as positive control. 0,100 ml spore suspension ($1,6 \times 10^6$ spores/0,1 ml) is diluted in 9,90 ml 40% ethanol 3 times to give 1,6 spores/0,1ml in the final suspension, which corresponds to 5 spores/1,6 spores = $3,125 \times 0,1 \text{ ml} = 0,3125 \text{ ml}$ spore suspension to each contact sheet (24 cm^2).

40

Class B: Spore suspensions containing 25 spores/63 μL is used as positive control. 0,100 ml spore suspension ($1,6 \times 10^6$ spores/0,1 ml) is diluted in 19,90 ml 40% ethanol 2 times to give 40 spores/0,1ml in the final suspension, which corresponds to 25 spores/40 spores = $0,625 \times 0,1 \text{ ml} = 0,0625 \text{ ml}$ spore suspension to each contact sheet (24 cm^2).

45

Negative control

Sterile ethanol is used as negative control

APPLICATION/32632DK01/PDK/LSH/11-12-02

Linearity

Three spore suspensions a) containing 5 spores, b) containing 25 spores and c) containing 50 spores will be used for the linearity study.

5

a) 0,100ml spore suspension ($1,6 \cdot 10^6$ spores/0,1 ml) is diluted in 9,90 ml 40% ethanol 3 times to give 1,6 spores/0,1 ml in the final suspension, which corresponds to 5 spores/1,6 spores = $3,125 \times 0,1 \text{ ml} = 0,3125 \text{ ml}$ spore suspension to each contact sheet (24 cm²).

10

b) 0,100 ml spore suspension ($1,6 \cdot 10^6$ spores/0,1 ml) is diluted in 19,90 ml 40% ethanol 2 times to give 40 spores/0,1ml in the final suspension, which corresponds to 25 spores/40 spores = $0,625 \times 0,1 \text{ ml} = 0,0625 \text{ ml}$ spore suspension to each contact sheet (24 cm²).

15

c) 0,100 ml spore suspension ($1,6 \cdot 10^6$ spores/0,1 ml) is diluted in 19,90 ml 40% ethanol 2 times to give 40 spores/0,1ml in the final suspension, which corresponds to 50 spores/40 spores = $1,25 \times 0,1 \text{ ml} = 0,125 \text{ ml}$ spore suspension to each contact sheet (24 cm²).

Sample preparation.

Sampling using swab technique.

20

- use gloves
- Spore suspension is applied to each stainless steel sheet
- Ethanol from the spore suspension on the stainless steel sheets is allowed to evaporate
- 1.0 ml alcalase solution is diluted in 90 ml saltwater peptone solution.
- 25 - put stomacher bag in matching stomacher bag clip
- swab the stainless steel sheets horizontally from left to right covering the 24 cm² contact area using a swab moistened with saltwater-peptone solution.
- swab the stainless steel sheets horizontally from top to bottom covering the 24 cm² contact area using a swab moistened with saltwater-peptone solution.
- 30 - swab the stainless steel sheets horizontally from left to right covering the 24 cm² contact area using a dry swab.
- swab the stainless steel sheets horizontally from top to bottom covering the 24 cm² contact area using a dry swab.
- place the swabs in stomacher bags.
- 35 - add the diluted alcalase solution to each stomacher bag.
- homogenize the stomacher bags using a stomacher.
- incubate the stomacher bags at 36C until the swabs have been dissolved (for at least 1 hour and maximum 2,5 hours).
- extraction of the spores is carried out using membrane filtration

40

Sampling using Count-Tact applicator

- use gloves
- spore suspension is applied to each stainless steel sheet
- 45 - ethanol from the spore suspension on the stainless steel sheets is allowed to evaporate
- Count-Tact applicator is used on the stainless steel plates

18

Negative controls are also added to the stainless steel sheets. Positive control samples are not added to stainless steel sheets, but are added directly to the Count-Tact application plates.

Incubation

- 5 All samples are incubated at $55\text{C} \pm 2\text{C}$ for at least 1 day to a maximum of 7 days.

Cleaning of surface material and templates.

Surface material and templates should be cleaned after each use and re-use:

- rinse the surface by wiping with laboratory paper that has been soaked in ethanol
- 10 - sterilization by autoclave at 121C for not less than 20 minutes

Calculations

Recovery% = $\text{sample (CFU)} * 100 / \text{added spores}$, where

sample = number of spores from Accuracy Test Solutions and

- 15 Added = number of spores in Accuracy Control Samples

Validation procedures

Recovery/Precision/Accuracy

20

Recovery/Precision and Accuracy is investigated using stainless steel and performed in accordance with the methods outlined above. This test will be used for the calculation of the microbiological recovery from the test samples. Batches of reagents and equipment will not be altered since the change in these parameters is estimated to have little or no influence on the

- 25 final test results.

Procedure

One analyst will contaminate six stainless steel plates with approx. 5 and 25 bacillus spores each.

- 30 Each stainless steel sheet is swabbed quadruplicate or sampled using the Count-Tact applicator. The samples are incubated at $55 \pm 2\text{C}$ for at least 1 day to a maximum of 7 days and counted. The tests will be carried out in accordance with the method described above.

Recovery/Accuracy

- 35 Recovery should be higher than 20% compared to positive controls.

The lowest recovery sets the limit (the lowest average recovery at any given spore concentration - 5 to 25 spores/ 24 cm^2).

Precision

- 40 - calculate mean and % RSD on replicate analysis
- % RSD should be less than 10%

Linearity

- 45 Procedure

19

Prepare six identical samples using the spore suspensions a), b) and c) defined above. The samples will be used to establish recovery efficiencies for these levels of spores and used for the subsequent linear regression.

5 Evaluation

- recovery will be calculated for each level of spores and regression will be made from 5 to 50 spores.
 - calculate the regression parameters of the standard curves and report slope, intercept and correlation coefficient of each curve.
- 10 - the correlation coefficient should meet the criterial of $r^2 > 0,940$.

Intermediate precision

Procedure

- 15 Same as for precision described above on three following days, using three different analysts.

Evaluation

- calculate recovery of spores
 - calculate mean and % RSD on replicate analysis for each day
- 20 - calculate mean and % RSD using all days (intermediate precision day 1, day 2 and day 3)
- % RSD should be less than 15%

Example 8. Validation of sampling from polished stainless steel.

25

Purpose

To validate the methods used for microbiological sampling from polished stainless steel that has been cleaned with 70% isopropanol, referring to the validation protocol of Example 7.

30 Introduction.

The validation study is carried out in order to define the accuracy, precision and linearity of methods used for microbiological sampling from surfaces and to estimate the recovery efficiency of micro-organisms from the type of material used in the test, using the described methods of sampling.

35

In the present study the sampling was performed from polished stainless steel cleaned with 70% isopropanol. The rationale for sampling is that this is a common material used for equipment production. The method tested were sampling by swab technique and sampling by count-tact technique.

40

The sampling was carried out on surfaces with an applied number of bacterial spores equal to the USP, NF guideline of class 10,000 production equipment and the EU-GMP guideline for microbiological purity of class 10,000 production equipment.

45

Results

Recovery

Calculation of the recovery using the swab technique (Table 1) showed variations between analysts when applying 5 spores (40% to 175%) and variations when applying 25 spores (73% to 105%). The average recovery for all analysts using the swab technique were calculated to 80% when applying 5 spores and 91% when applying 25 spores.

Calculation of recovery using the count-tact technique (Table 2) showed variations between analysts when applying 5 spores (54% to 88%) and variations when applying 25 spores (36% to 57%). The average recovery for all analysts using the count-tact technique were calculated to 74% when applying 5 spores and 45% when applying 25 spores.

Precision

Calculations of relative standard deviation in percent (RSD %), when using the swab technique (Table 1) showed variations between analysts (34% to 70%) when applying 5 spores and variation (30% to 57%) when applying 25 spores. Using the count-tact technique (Table 2) RSD% was between 41% and 90% when applying 5 spores and between 20% and 65% when applying 25 spores.

Intermediate precision

Intermediate precision using the swab technique (Table 1) was 64% when applying 5 spores and 43% when applying 25 spores. Intermediate precision using the count-tact technique (Table 2) was 77% when applying 5 spores and 54% when applying 25 spores.

Table 1. Sampling results using the swab technique

Spores applied (calculated, not actual) ¹⁾		Assay (spores/24 cm ²)		
		5	25	50
Mean (cfu/24 cm ²)	Analyst 1 ²⁾	5	21	2)
Precision (RSD%)		42	57	
Recovery (%)		175	105	
Mean (cfu/24 cm ²)	Analyst 2 ³⁾	3	15	2)
Precision (RSD%)		70	30	
Recovery (%)		59	73	
Mean (cfu/24 cm ²)	Analyst 3 ³⁾	2	17	29
Precision (RSD%)		34	32	39
Recovery (%)		40	94	95
Mean (cfu/24 cm ²)	Analyst 1,2,3	3	18	2)
Precision (RSD%)		64	43	
Recovery (%)		80	91	

¹⁾The number of spores applied was calculated to be 5, 25 or 50. The actual amount of spores applied was used for calculating recoveries.

²⁾One analyst only carried out sampling from surfaces with 50 spores applied by the sole purpose of investigating linearity

³⁾Each analyst completed the sampling and analysis in six replicates.

21

Linearity from 5 to 50 spores:

Correlation coefficient (R) = 0,9995

$R^2 = 0,9990$

5 Intercept = -2,3

Slope = 1,0

Linearity

10 Linearity was calculated by regression between applied spore concentrations of 5, 25 and 50 spores by one analyst. The defined level of acceptance was a $R^2 > 0,9400$. The calculated R^2 was 0,9990 when using the swab technique and 0,9989 when using the count-tact technique.

Table 2. Sampling results using the count-tact technique

Spores applied (calculated, not actual) ¹⁾		Assay (spores/24 cm ²)		
		5	25	50
Mean (cfu/24 cm ²)	Analyst 1 ³⁾	4	14	20
Precision (RSD%)		70	20	24
Recovery (%)		88	45	40
Mean (cfu/24 cm ²)	Analyst 2 ³⁾	2	7	2)
Precision (RSD%)		41	44	
Recovery (%)		79	36	
Mean (cfu/24 cm ²)	Analyst 3 ³⁾	2	13	2)
Precision (RSD%)		90	65	
Recovery (%)		54	57	
Mean (cfu/24 cm ²)	Analyst 1,2,3	3	11	2)
Precision (RSD%)		77	54	
Recovery (%)		74	45	

15 ¹⁾The number of spores applied was calculated to be 5, 25 or 50. The actual amount of spores applied was used for calculating recoveries.

²⁾One analyst only carried out sampling from surfaces with 50 spores applied by the sole purpose of investigating linearity

³⁾Each analyst completed the sampling and analysis in six replicates.

20 Linearity from 5 to 50 spores:

Correlation coefficient (R) = 0,9989

$R^2 = 0,9979$

Intercept = -2,4

Slope = 0,4

25

Discussion

Recovery

- The level of recovery of microorganisms from surfaces is critical when complying with USP/NF guidelines and EU-GMP guidelines. The bacterial recovery using both the swab technique and the count-tact technique as described in the protocol was better than anticipated when recovering microorganisms from the samples with a known microbiological contamination equal to the USP/NF guideline and the EU-GMP guideline. However, the recovery efficiency varied depending on the level of applied spores. Thus, the lowest recovery efficiency should be used.
- 10 The average (for all analysts) lowest recovery was 80% for the swab technique and 45% for the count-tact technique. For the chosen sampling method used on stainless steel this recovery should be used to estimate the actual amount of microorganism on the surface.

Precision

- 15 The relative standard deviation (%RSD) on replicate analysis (for a single analyst) was larger than first anticipated. The comparison of the %RSD for the positive controls and the assay however showed that high %RSD is mainly caused by the analysis of the samples and not by the use of count-tact or swab technique. It should have been expected not to achieve a %RSD
- 20 corresponding to chemical analysis, since it is generally known that %RSD is much larger for microbiological analysis than chemical analysis.

Intermediate precision

- 25 The relative standard deviation (%RSD) on replicate analysis (between analysts) was larger than first anticipated. The comparison of the %RSD for the positive controls and the assay however showed that the high %RSD is mainly caused by the analysis of the samples and not by the swab or count-tact techniques used. It should have been expected not to achieve a %RSD corresponding to chemical analysis, since it is generally known that %RSD is much larger for microbiological analysis than chemical analysis.
- 30 The average %RSD calculated for the lowest recoveries given above was 64% for the swab technique and 54% for the count-tact method. To correct for the large variations on the analysis the %RSD should be taken into account when estimating the actual number of microorganisms on a surface.

Linearity

- 35 Both methods of sampling give a high degree of linearity at the spore concentrations used in the present validation study. Correlation was better than anticipated.

Conclusions

- 40 The results show that the methods described above can be used for sampling of microorganisms from polished stainless steel in an overall satisfactory way. The level of recovery of microorganisms from surfaces is critical when complying with USP/NF guidelines and EU-GMP guidelines. The bacterial recovery using both the swab technique and the count-tact applicator as described in the protocol was better than anticipated when recovering microorganisms from
- 45 samples with a known microbiological contamination equal to the USP/NF guideline and the EU-GMP guideline.

CLAIMS

1. A device comprising
 - i) a gelatin-based sponge; and
 - 5 ii) a supportwherein said sponge comes into contact with or is attached to said support.
2. The device according to claim 1 wherein said support is made from materials selected from the group consisting of wood, plastics, rubber, and natural or synthetic polymeric material.
- 10 3. The device according to claim 1 wherein said support is of the form selected from the group consisting of a disc, a cube, a block, a sphere, a crucible, a coating and a handle.
4. The device according to claim 1 wherein said support is in the form of a handle.
- 15 5. The device according to claim 4 wherein said handle is in the form of a stick.
6. The device according to claim 1 wherein the gelatin-based sponge is based on organic or synthetic gelatin, such as gelatin from the group consisting of mammals, marine mammals,
- 20 porcine, bovine, fish, crayfish or vegetables.
7. The device according to claim 1 wherein the gelatin-based sponge is based on gelatin of porcine origin.
- 25 8. The device according to claim 1, wherein the gelatin-based sponge has a reconfirmation of no more than 10 seconds, typically no more than 5 sec.
9. The device according to claim 1, wherein the gelatin-based sponge has a water absorption capacity of at least 30 g/g, typically at least 40 g/g.
- 30 10. The device according to claim 1, wherein the gelatin-based sponge is in the form of a powder.
11. The device according to claim 10, wherein the powder has a particle size in the range of about 1 μ m to about 1 mm, such as about 5 μ m to about 0.5 mm, such as about 10 μ m to about 0.1 mm.
- 35 12. The device according to claim 1, wherein the powder is incorporated into a binding material.
- 40 13. The device according to claim 12, wherein the binding material is a swab, such as a cotton-based swab.
14. The device according to claim 1, wherein the gelatin-based sponge is constituted by at least
- 45 50% gelatin, such as at least 60%, such as at least 70%, typically at least 75% gelatin, preferably at least 80%, more preferably at least 85% gelatin, such as at least 90% gelatin,

suitably at least 95% gelatin, most preferably selected from at least 96%, 97%, 98%, 99% gelatin, based on the dried weight of the sponge.

15. A device comprising

- 5 i) a gelatin-based sponge; and
ii) an agent selected from the group consisting of a neutral diluent, an anti-microbial agent and a dispersion agent.

16. The device according to claim 15, wherein said neutral diluent is selected from the group
10 consisting of saline, saline peptone, buffered saline peptone, Ringer solution and an organic or inorganic buffer.

17. The device according to claim 15, wherein said gelatin-based sponge comes into contact
15 with or is attached to a support.

18. The device according to claim 17 wherein said support is of the form selected from the group consisting of a disc, a cube, a block, a sphere, a crucible, a coating and a handle.

19. The device according to claim 17 wherein said support is in the form of a handle.

- 20 20. The device according to claim 19 wherein said handle is in the form of a stick.

21. The device according to claim 15 wherein the gelatin-based sponge is based on organic or synthetic gelatin, such as gelatin from the group consisting of mammals, marine mammals,
25 porcine, bovine, fish, crayfish or vegetables.

22. The device according to claim 15 wherein the gelatin-based sponge is based on gelatin of porcine origin.

- 30 23. The device according to claim 15, wherein the gelatin-based sponge has a reconfirmation of no more than 10 seconds, typically no more than 5 sec.

24. The device according to claim 15, wherein the gelatin-based sponge has a water absorption capacity of at least 30 g/g, typically at least 40g/g.

- 35 25. The device according to claim 15, wherein the gelatin-based sponge is in the form of a powder.

26. The device according to claim 25, wherein the powder has a particle size in the range of
40 about 1 μ m to about 1 mm, such as about 5 μ m to about 0.5 mm, such as about 10 μ m to about 0.1 mm.

27. The device according to claim 25, wherein the powder is incorporated into a binding material.

- 45 28. The device according to claim 27, wherein the binding material is a swab, such as a cotton-based swab.

29. The device according to claim 15, wherein the gelatin-based sponge is constituted by at least 50% gelatin, such as at least 60%, such as at least 70%, typically at least 75% gelatin, preferably at least 80%, more preferably at least 85% gelatin, such as at least 90% gelatin, suitably at least 95% gelatin, most preferably selected from at least 96%, 97%, 98%, 99% gelatin, based on the dried weight of the sponge.
30. Use of a gelatin-based sponge for collection of a target from a collection medium comprising making contact between the gelatin-based sponge and the target.
31. The use according to claim 30, wherein the collection medium is selected from the group consisting of a solid or semi-solid surface, a liquid, a gas and combinations thereof.
32. The use according to claim 30, wherein the target is selected from the group consisting of a micro-organism, a mammalian cell and a molecular target.
33. The use according to claim 32, wherein the molecular target is a nucleotide, a nucleic acid or a detergent.
34. The use according to claim 33, wherein the nucleotide is a purine- or a pyrimidine-containing nucleotide, preferably ATP.
35. The use according to claim 32, wherein the micro-organism is selected from the group consisting of bacteria, archaea, bacterial spores, yeast, fungi and viruses.
36. The use according to any one of claims 30-35 further comprising the step of transfer of the target from the sponge to a first transfer medium.
37. The use according to claim 36, wherein the transfer comprises the digestion of the gelatin-based sponge.
38. The use according to claim 36, wherein the transfer comprises the mechanical transfer of target from the gelatin-based sponge to a second transfer medium.
39. The use according to claim 36, wherein the transfer comprises the washing of target from the gelatin-based sponge.
40. The use according to claim 37, wherein the digestion comprises the use of an agent selected from the group consisting of an enzyme, a mineral acid, a carboxylic acid, a base and combinations thereof.
41. The use according to claim 40, wherein the enzyme is a protease.
42. The use according to claim 41, wherein the protease is alcalase or pepsin.
43. The use according to any of the claims 36-42, further comprising the step of extraction of the target by membrane filtration.

44. The use according to claim 30, wherein the gelatin-based sponge has a reconfirmation of no more than 10 seconds, typically no more than 5 sec.
- 5 45. The use according to claim 30, wherein the gelatin-based sponge has a water absorption capacity of at least 30 g/g, typically at least 40g/g.
46. The use according to claim 30, wherein the gelatin-based sponge is in the form of a powder.
- 10 47. The use according to claim 46, wherein the powder has a particle size in the range of about 1 μ m to about 1 mm, such as about 5 μ m to about 0.5 mm, such as about 10 μ m to about 0.1 mm.
- 15 48. The use according to claim 46, wherein the powder is incorporated into a binding material.
49. The use according to claim 48, wherein the binding material is a swab, such as a cotton-based swab.
- 20 50. The use according to claim 30, wherein the gelatin-based sponge is constituted by at least 50% gelatin, such as at least 60%, such as at least 70%, typically at least 75% gelatin, preferably at least 80%, more preferably at least 85% gelatin, such as at least 90% gelatin, suitably at least 95% gelatin, most preferably selected from at least 96%, 97%, 98%, 99% gelatin, based on the dried weight of the sponge.
- 25 51. A method of lowering the amount of a target in a sample area comprising making contact between a gelatin-based sponge and at least a portion of said sample area, such that the target adheres to the sponge.
- 30 52. The method according to claim 51, wherein the collection medium is selected from the group consisting of a solid or semi-solid surface, a liquid, a gas and combinations thereof.
53. The method according to claim 51, wherein the target is selected from the group consisting of a micro-organism, a mammalian cell and a molecular target.
- 35 54. The method according to claim 51, wherein the molecular target is a nucleotide, a nucleic acid or a detergent.
55. The method according to claim 54, wherein the nucleotide is a purine- or a pyrimidine-
40 containing nucleotide, preferably ATP.
56. The method according to claim 53, wherein the micro-organism is selected from the group consisting of bacteria, archaea, bacterial spores, yeast, fungi and viruses.
- 45 57. The method according to any one of claims 51-56 further comprising the step of transfer of the target from the sponge to a first transfer medium.

58. The method according to claim 57, wherein the transfer comprises the digestion of the gelatin-based sponge.
59. The method according to claim 57, wherein the transfer comprises the mechanical transfer
5 of the target from the gelatin-based sponge to a second transfer medium.
60. The method according to claim 57, wherein the transfer comprises washing of a target from the gelatin-based sponge.
- 10 61. The method according to claim 51, wherein the gelatin-based sponge has a reconfirmation of no more than 10 seconds, typically no more than 5 sec, as determined by the method of Example 1.
62. The method according to claim 51, wherein the gelatin-based sponge has a water
15 absorption capacity of at least 30 g/g, typically at least 40g/g, as determined by the method of Example 3.
63. The method according to claim 51, wherein the gelatin-based sponge is in the form of a powder.
20
64. The method according to claim 63, wherein the powder has a particle size in the range of about 1 μ m to about 1 mm, such as about 5 μ m to about 0.5 mm, such as about 10 μ m to about 0.1 mm.
- 25 65. The method according to claim 63, wherein the powder is incorporated into a binding material.
66. The method according to claim 65, wherein the binding material is a swab, such as a cotton-based swab.
30
67. The method according to claim 51, wherein the gelatin-based sponge is constituted by at least 50% gelatin, such as at least 60%, such as at least 70%, typically at least 75% gelatin, preferably at least 80%, more preferably at least 85% gelatin, such as at least 90% gelatin, suitably at least 95% gelatin, most preferably selected from at least 96%, 97%, 98%, 99%
35 gelatin, based on the dried weight of the sponge.
68. The method according to any one of claims 51-67, wherein the gelatin-based sponge is presterilized.
- 40 69. The method according to any one of claims 51-67, wherein the gelatin-based sponge further comprises an agent selected from the group consisting of a neutral diluent, an anti-microbial agent, a disinfecting agent and a dispersion agent.
70. The method according to claim 69, wherein the anti-microbial or disinfecting agent is an
45 alcohol.

71. A method of qualitatively or quantitatively sampling an area for content of a target comprising the use of a gelatin-based sponge and the steps of
i) wet sampling using the gelatin-based sponge; and/or
ii) dry-sampling using the gelatin-based sponge.
- 5
72. The method according to claim 71, wherein the collection medium is selected from the group consisting of a solid or semi-solid surface, a liquid, a gas and combinations thereof.
73. The method according to claim 71, wherein the target is selected from the group consisting
10 of a micro-organism, a mammalian cell and a molecular target.
74. The method according to claim 73, wherein the molecular target is a nucleotide, a nucleic acid or a detergent.
- 15
75. The method according to claim 74, wherein the nucleotide is a purine- or a pyrimidine-containing nucleotide, preferably ATP.
76. The method according to claim 73, wherein the micro-organism is selected from the group consisting of bacteria, archaea, bacterial spores, yeast, fungi and viruses.
- 20
77. The method according to any one of claims 71-76 further comprising the step of transfer of the target from the sponge to a first transfer medium.
78. The method according to claim 77, wherein the transfer comprises the digestion of the
25 gelatin-based sponge.
79. The method according to claim 77, wherein the transfer comprises the mechanical transfer of target from the gelatin-based sponge to a second transfer medium.
- 30
80. The method according to claim 77, wherein the transfer comprises washing of a target from the gelatin-based sponge.
81. The method according to claim 78, wherein the digestion comprises the use of an agent selected from the group consisting of an enzyme, a mineral acid, a carboxylic acid, a base and
35 combinations thereof.
82. The method according to claim 78, wherein the enzyme is a protease.
83. The method according to claim 82, wherein the protease is alcalase or pepsin.
- 40
84. The method according to any of the claims 77-83, further comprising the step of extraction of the target by membrane filtration.
85. The method according to claim 71, wherein the gelatin-based sponge has a reconfirmation
45 of no more than 10 seconds, typically no more than 5 sec.

86. The method according to claim 71, wherein the gelatin-based sponge has a water absorption capacity of at least 30 g/g, typically at least 40g/g, as determined by the method of Example 3.
- 5 87. The method according to claim 71, wherein the gelatin-based sponge is in the form of a powder.
88. The method according to claim 87, wherein the powder has a particle size in the range of about 1 μ m to about 1 mm, such as about 5 μ m to about 0.5 mm, such as about 10 μ m to
10 about 0.1 mm.
89. The method according to claim 87, wherein the powder is incorporated into a binding material.
- 15 90. The method according to claim 89, wherein the binding material is a swab, such as a cotton-based swab.
91. The method according to claim 71, wherein the gelatin-based sponge is constituted by at least 50% gelatin, such as at least 60%, such as at least 70%, typically at least 75% gelatin, preferably at least 80%, more preferably at least 85% gelatin, such as at least 90% gelatin, suitably at least 95% gelatin, most preferably selected from at least 96%, 97%, 98%, 99%
20 gelatin, based on the dried weight of the sponge.
92. A method for culturing micro-organisms or mammalian cells comprising adhering the cells
25 to a gelatin-based sponge and culturing the cells in a growth medium.
93. The method according to claim 92, wherein the growth medium is added to the gelatin-based sponge.
- 30 94. The method according to claim 92, wherein the gelatin-based sponge is incubated in a liquid growth medium.
95. The method according to claim 92, further comprising the step of digesting the gelatin-based sponge.
35
96. The method according to claim 92, wherein the digestion comprises the use of an agent selected from the group consisting of an enzyme, a mineral acid, a carboxylic acid, a base and combinations thereof.
- 40 97. The method according to claim 96, wherein the enzyme is a protease.
98. The method according to claim 97, wherein the protease is alcalase or pepsin.
99. The method according to claim 92, wherein the gelatin-based sponge has a reconfirmation
45 of no more than 10 seconds, typically no more than 5 sec.

30

100. The method according to claim 92, wherein the gelatin-based sponge has a water absorption capacity of at least 30 g/g, typically at least 40g/g.

101. The method according to claim 92, wherein the gelatin-based sponge is in the form of a powder.

102. The method according to claim 101, wherein the powder has a particle size in the range of about 1 μ m to about 1 mm, such as about 5 μ m to about 0.5 mm, such as about 10 μ m to about 0.1 mm.

10

103. The method according to claim 101, wherein the powder is incorporated into a binding material.

104. The method according to claim 103, wherein the binding material is a swab, such as a cotton-based swab.

105. The method according to claim 92, wherein the gelatin-based sponge is constituted by at least 50% gelatin, such as at least 60%, such as at least 70%, typically at least 75% gelatin, preferably at least 80%, more preferably at least 85% gelatin, such as at least 90% gelatin, suitably at least 95% gelatin, most preferably selected from at least 96%, 97%, 98%, 99% gelatin, based on the dried weight of the sponge.

20